## POISON ANTIDO

Utah Poison Control Center 585 Komas Drive Salt Lake City, Utah 84108 1-800-222-1222

## **CARBON MONOXIDE POISONING**

Carbon monoxide (CO) is a colorless, odorless, and tasteless, and can be deadly in certain concentrations. When an individual is exposed to carbon monoxide, the ability for blood to carry oxygen throughout the body is lessened; body tissues and organs can suffocate.

CO can escape into a home through a furnace that isn't working correctly, a wood burning stove, water heater, gas range, or a fireplace.

There is a way to prevent CO poisoning. Make sure to install and use high quality carbon monoxide detectors. CO detectors should be placed near bedrooms and if your home has more than one level, at least one detector per level in the home.

Before the start of the winter season have the furnace and heating system in your home inspected, warm up an automobile in an open garage or outside and have your chimney and flu cleaned professionally.

How can you tell if you have been potentially poisoned by CO? Carbon Monoxide poisoning symptoms can be easily confused with flu symptoms but are much more serious. Symptoms include:

- Headache
- Fatigue
- Dizziness
- Nausea
- Irregular Breathing
- Confusion
- Unconsciousness



If you think someone has been poisoned by CO, call the Poison Control Center immediately and take the person to a well ventilated area. If the person is unconscious or having difficulty breathing, call 9-1-1. Whether it's a poison emergency or a poisoning question, please do not hesitate to call us at 1-800-222-1222.

## **INSIDE THIS ISSUE**

Carbon Monoxide Poisoning What's in the Garage Cough and Cold Preparations

1

WHAT'S IN THE GARAGE?

The winter months bring different maintenance concerns for automobiles. Antifreeze and windshield washer fluid are necessary to keep cars functioning and safe. Here are some things to keep in mind when using such products:

- Please keep all types of chemicals in their original containers.
- Make sure to store products in the garage high up and in a locked cabinet.
- Brightly colored chemicals are attractive to young children.
- Most poisonings occur while the product is being used.

If someone does get poisoned please have the product with you and call the Poison Control Center right away!

## **COUGH AND COLD PREPARATIONS**

The winter months bring the flu, colds, aches, and pains. This is a good time to make sure that the cough and cold preparations you are taking are safe to take along with over the counter (OTC) medications or prescriptions drugs. Here are a few things to think about when considering medicines for your coughs and colds.

- The active ingredients acetaminophen and ibuprofen are used in many OTC medicines. Some products include: Tylenol®, TheraFlu®, NyQuil®, Advil®, and Motrin®.
- To help prevent an overdose, the drug facts label will help you know active ingredients in medicines that might also be in medicines you are taking.

 Review warnings on all prescription and OTC medications.

If you think a potentially poisonous drug interaction or overdose has occurred, please do not hesitate to contact the Poison Control Center.

Drug Facts		
Criterpheniramine maleate 2 mg	iblet)	
Chlorphenizanine maleste 2 mg  Uses temporarily relieves those symptimal sneezing manny nose maleste with the second manny nose maleste with the second manny nose malesters.	English Control of Con	Purpos
aneering a nummy none as sympto	Cetts clue to hely forms or -0	Anthistamin
Uses temporarily relieves those sympo- sneezing Interry nose I fally, will Warnings	Mary eyes # Italy thouse	piratory afernies
Ask a doctor before use if you have		
troucke unnation due to	At employees	
glaucoma a breathing problem such a trouble uninstrop due to an entarped prob	state plant or cryonic bronchis	
	Target and the same and the sam	
When using this product	you are taking tranquitzers or sectative	
		10
alcohol, secialises, and transpulsave may a alcohol, secialises, and transpulsave may a secareful when driving a motor whice or exceptibly may occur, especially in children to present the second	d alcoholic drinka	
months when driving a motor vehicle or	CONTRACTOR CHORESTONES	
If present in children	nacranery	
If pregnant or breast-feeding, ask a health, Keep out of reach of children, in case of ov Center right away.  Directions	and I	
Center right away	proressional before use.	_
Directions	per medical help or contact a F	Minns di
- rections		CHARLI CIGINION
adults and children 12 years and over	No. of the last of	
1 and over	take 2 tablets every 4 to 6 ho	
Children & union to	not more than 12 tablets in 24	ung:
	tring 1 to the company of the	
children 6 years to under 17 years		4
children under 6 years	take 1 tablet every 4 to 6 hour not more than 6 tablets in	
children under 6 years		
other ender 6 years	ask a doctor	Nous
	ask a doctor	Nous